

These select priority health indicators from <u>www.CityHealthDashboard.com</u> for Springfield, MO and Lexington, KY provide an overview of health outcomes and the community factors that contribute to the overall health of the community, as well as illuminate opportunities for improvement.

Comparing Springfield and Lexington		
	Springfield, MO	Lexington, KY
Overall Score	51.8	61.3
The green bars are a calculated score from 1 (worst) to 100 (best) among the 500 largest cities in the U.S. with a collection of data—not all indicators which make up that score are included in this document.		
Clinical Care	42.1	68.2
Prenatal care	70%	80%
	Births for which pre	enatal care began in the first trimester
Health Behavior	52.0	57.6
Physical inactivity	28%	25%
· · · ·	No leisure-time physical act	ivity in past month among adults $\geq$ 18
Smoking	25%	21%
	·	Current smoking among adults ≥ 18
Health Outcomes	53.3	59.4
Cardiovascular disease deaths	227.0	169.3
	·	Per 100,000 population
Frequent mental distress	16%	13%
Mer	ntal health not good for $\geq$ 14 days durin	ig the past 30 days among adults ≥18
Life expectancy	77.6	78.0
		Average life expectancy at birth
Obesity	32%	32%
		Among adults ≥ 18
Opioid overdose deaths	17.8	27.4
		Per 100,000 population
Physical Environment	45.5	45.0
Park access	46%	52%
	Population living w	ithin a 10-minute walk of green space
Limited access to healthy foods	73%	71%
· · · · ·	half-mile from the nearest supermark	
Walkability	38.4	34.2
5	borhood amenities accessible by walki	· ·
Social and Economic Factors	57.4	72.1
Children in Poverty	30%	23%
Children living in households on or above the federal poverty line		
High school graduation	88%	88%
	· · · ·	within four years of entering 9th grade
Income Inequality	-29.9	-9.2
	e extremes of the national income distr	•
Third-grade reading proficiency	52%	55%
Th	nird-graders who score "proficient" or a	bove in reading on standardized tests